

UREA CYCLE DISORDERS DON'T AFFECT JUST NEONATES

confusion aggressiveness vomiting learning disabilities urea cycle disorders neurodevelopmental delay acute encephalopathy stroke-like episodes late onset poor appetite confusion hyperammonemia hyperammonemia mental retardation dizziness protein aversion NAGS deficiency hyperactivity confusion vomiting somnolence ataxia failure to thrive Seizures lethargy coma aggressiveness mood alteration ataxia liver failure seizures lethargy mental retardation aggressiveness hyperammonemia Urea Cycle Disorders coma

Hyperammonemia due to NAGS deficiency may occur at any time of life

INDICATIONS AND USAGE

CARBAGLU® (carglumic acid) is a Carbamoyl Phosphate Synthetase 1 (CPS1) activator indicated as:

- Adjunctive therapy in pediatric and adult patients for the treatment of acute hyperammonemia due to the deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS). During acute hyperammonemic episodes, concomitant administration of CARBAGLU with other ammonia lowering therapies, such as alternate pathway medications, hemodialysis, and dietary protein restriction, is recommended.
- Maintenance therapy in pediatric and adult patients for the treatment of chronic hyperammonemia due to the deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS). During maintenance therapy, the concomitant use of other ammonia lowering therapies and protein restriction may be needed based on plasma ammonia levels.


IMPORTANT SAFETY INFORMATION

- Most common adverse reactions (>9%) are: vomiting, abdominal pain, pyrexia, tonsillitis, anemia, diarrhea, ear infection, infections, nasopharyngitis, hemoglobin decreased, and headache.

Carbaglu[®]
carglumic acid

UREA CYCLE DISORDERS DON'T AFFECT JUST NEONATES...DIAGNOSING LATE-ONSET NAGS DEFICIENCY PATIENTS

UREA CYCLE AND DEFECTS



- In humans, nitrogen is produced by catabolism of proteins and other nitrogen-containing molecules
- The urea cycle converts waste nitrogen into urea
- Deficiency of enzymes in the urea cycle may lead to a build-up of ammonia, which is neurotoxic

NAGS: N-acetylglutamate synthase

CPS1: carbamoyl phosphatase 1

OTC: ornithine transcarbamylase

ASS: argininosuccinate synthetase

ASL: argininosuccinate lyase

ARG1: arginase


Interruption of the urea cycle causes hyperammonemia which, if untreated, may result in coma and death

The mortality rate during hyperammonemic crises can reach 10%

IMPORTANT SAFETY INFORMATION

- **Hyperammonemia:** Monitor plasma ammonia level during treatment. Prolonged exposure to elevated plasma ammonia level can result in brain injury or death. Prompt use of all therapies necessary to reduce plasma ammonia level is essential.
- **Pregnancy:** No human data; decreased survival and growth in animal offspring.
- **Nursing Mothers:** Breastfeeding is not recommended while taking CARBAGLU.

DIAGNOSTIC ALGORITHM FOR HYPERAMMONEMIA IN UCDs



Adapted from Häberle J, et al. Suggested guidelines for the diagnosis and management of urea cycle disorders. *Orphanet J Rare Dis.* 2012;7:32.

Genetic testing is virtually the only method that can confirm a diagnosis of NAGS deficiency. For more information consult the National Center for Biotechnology Information (NCBI) Genetic Testing Registry at www.ncbi.nlm.nih.gov/gtr

IMPORTANT SAFETY INFORMATION

- **Hyperammonemia:** Monitor plasma ammonia level during treatment. Prolonged exposure to elevated plasma ammonia level can result in brain injury or death. Prompt use of all therapies necessary to reduce plasma ammonia level is essential.

NAGS DEFICIENCY

- NAGS deficiency is a rare autosomal recessive urea cycle disorder
- The classic presentation of urea cycle disorders including primary NAGS deficiency is high levels of ammonia in the neonatal period
- Partial/milder enzyme deficiency may permit an individual to function normally until a stressor triggers a hyperammonemic crisis:
 - Valproic acid | Heart-lung transplant | Parenteral nutrition with high nitrogen intake
 - Post-partum stress | Short bowel and kidney disease | Gastrointestinal bleeding

NAGS DEFICIENCY IN LATE-ONSET PATIENTS: CLINICAL PRESENTATION

FINDINGS IN REPORTED CASES OF CONFIRMED NAGS DEFICIENCY

GASTROINTESTINAL PRESENTATIONS	NEUROLOGICAL PRESENTATIONS	PSYCHIATRIC PRESENTATIONS
<ul style="list-style-type: none">- Nausea, vomiting	<ul style="list-style-type: none">- Disorientation- Dilated pupils with delayed response to light- Exaggerated deep tendon reflexes- Generalized cerebral edema- Unresponsiveness to stimuli- Seizures- Atactic gait	<ul style="list-style-type: none">- Aggressive attitude- Restlessness- Episodic altered mental status- Confusion- Post-operative combativeness, combative behavior

BLOOD AMMONIA 113-4781 $\mu\text{mol/L}$ *

* Normal blood ammonia level in adults: 11-32 $\mu\text{mol/L}$

DIAGNOSTIC CLUES AND MANAGEMENT

- Plasma ammonia should be measured promptly in all patients with unexplained encephalopathy, including cyclical manifestations, to identify possible underlying metabolic disorders
- Prompt diagnosis and initiation of treatment are necessary to avoid potentially poor neurological outcomes
- Patients should be managed by physicians and medical teams experienced in metabolic disorders
- The mainstay of ongoing management of NAGS deficiency is maintenance of plasma ammonia level in a normal range by avoiding catabolic stress and using carglumic acid

Carglumic acid is the treatment of choice in NAGS deficiency

IMPORTANT SAFETY INFORMATION

- Most common adverse reactions (>9%) are: vomiting, abdominal pain, pyrexia, tonsillitis, anemia, diarrhea, ear infection, infections, nasopharyngitis, hemoglobin decreased, and headache.
- **To report SUSPECTED ADVERSE REACTIONS, contact Recordati Rare Diseases Inc. at 1-888-575-8344, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.**

ILLUSTRATIVE CASE OF NAGS DEFICIENCY DIAGNOSED IN A 38-YEAR-OLD MALE

20-year history of fluctuating behavioral changes associated with nausea and vomiting

Past medical history

Symptomatic (behavioral changes, confusion) in early childhood and adulthood

Ammonia levels never checked before hospitalization

Negative family history

Clinical course... in the emergency room

Ongoing nausea, vomiting, headache

Stable vital signs

Behavioral disinhibition and fluctuating drowsiness

Impaired coordination

Normal cranial nerve examination and power testing

Mild spasticity on muscle tone assessment

Significant asterixis (flapping tremor)

Otherwise unremarkable general examination

... during hospital admission and follow-up

High blood ammonia levels
434 $\mu\text{mol/L}$

Mild respiratory alkalosis

Normal head/spine MRI

Continuous EEG monitoring:
generalized encephalopathy

Elevated glutamine level (1062 $\mu\text{mol/L}$),
normal citrulline and urine amino acid levels

Diagnosis

Molecular sequencing of the NAGS gene: compound heterozygote for E433G and IVS6+5 G>A, (both mutations) with residual NAGS expression that may explain the absence of neonatal hyperammonemia and delayed presentation in adult life

Medical management

Low protein diet

Initially sodium phenylbutyrate (200 mg TID*) and citrulline (50 mg/kg TID)

Switched to carglumic acid (1200 mg TID) with a more liberalized protein intake

Outcome

Ammonia level returned to normal (29-35 $\mu\text{mol/L}$)

No hyperammonemic crisis for 2 years

Markedly improved behavior; short term memory loss remains

* TID: three times a day

ORDERING CARBAGLU® (carglumic acid)

Exclusive Distribution Through Accredo

Representatives at Accredo Health Group, Inc. are committed to helping you and your patients through the ordering process. CARBAGLU is not available in retail pharmacies.

Patient Home Delivery

1. Download the CARBAGLU Prescription & Enrollment Form from the Accredo website (www.accredo.com).
2. Complete the CARBAGLU Prescription & Enrollment Form and fax it to the Accredo specialty pharmacy department at 1-888-454-8488.
3. A CARBAGLU specialty pharmacy representative may call to verify information and determine next steps. In some cases, prior authorization may be necessary.
4. Questions? Contact a CARBAGLU specialty pharmacy representative at 1-888-454-8860.

Hospital Orders – Emergency or Inpatient

1. Alert your hospital pharmacy that your patient requires CARBAGLU. Send an order for CARBAGLU to your hospital pharmacy and specify whether it is a STAT order (required in 6 hours or less*).
2. Ask your hospital pharmacy to call the wholesale department at Accredo at 1-877-900-9223 to place the order.
3. The wholesale department at Accredo will work with your hospital pharmacy to obtain payment information, establish shipping timelines, and verify the delivery address.
4. The Accredo specialty pharmacy team will follow up in 1-2 days to help set up patient home delivery of CARBAGLU, if applicable.

*Delivery time frame is weather dependent and is not guaranteed.

FINANCIAL ASSISTANCE PROGRAMS

For patients experiencing financial hardship, Recordati Rare Diseases Inc. supports a Patient Assistance Program and a Co-Pay Assistance Program, administered by Accredo. For more information, call: 1-888-454-8860.

For more information about CARBAGLU, visit: www.carbaglu.net

Please see enclosed Full Prescribing Information, including Instructions for Use.



CARBAGLU is a licensed trademark of Recordati Rare Diseases Inc.
© 2019 Recordati Rare Diseases Inc.
www.recordatirarediseases.com PP-CBGL-US-0131

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use CARBAGLU safely and effectively. See full prescribing information for CARBAGLU.

CARBAGLU® (carglumic acid) tablet for oral suspension

Initial U.S. Approval: 2010

RECENT MAJOR CHANGES

Dosage and Administration (2)	11/2017
Warnings and Precautions (5.1)	11/2017

INDICATIONS AND USAGE

CARBAGLU is a Carbamoyl Phosphate Synthetase 1 (CPS 1) activator indicated as:

- Adjunctive therapy for the treatment of acute hyperammonemia due to the deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS). (1.1)
- Maintenance therapy for the treatment of chronic hyperammonemia due to the deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS). (1.2)

DOSAGE AND ADMINISTRATION

Dosing (2.1):

- **Acute Hyperammonemia:** The recommended initial pediatric and adult dosage is 100 mg/kg/day to 250 mg/kg/day. Titrate based on plasma ammonia level and clinical symptoms.
- **Maintenance for chronic hyperammonemia:** The recommended pediatric and adult maintenance dosage is 10 mg/kg/day to 100 mg/kg/day. Titrate to target normal plasma ammonia level for age.
- Divide the total daily dose into two to four doses.

- Monitor plasma ammonia and adjust the dosage to maintain the level within the normal range for age.

Preparation and Administration (2.2):

- Disperse CARBAGLU tablets in water. Do not swallow whole or crushed.
- Take immediately before meals or feedings.
- For instructions on administration orally or through a nasogastric tube, see the full prescribing information.

DOSAGE FORMS AND STRENGTHS

Tablets for oral suspension: 200 mg, functionally scored (3)

CONTRAINDICATIONS

None. (4)

WARNINGS AND PRECAUTIONS

Hyperammonemia: Monitor plasma ammonia level during treatment. Prolonged exposure to elevated plasma ammonia level can result in brain injury or death. Prompt use of all therapies necessary to reduce plasma ammonia level is essential. (5.1)

ADVERSE REACTIONS

Most common adverse reactions (>9%) are: vomiting, abdominal pain, pyrexia, tonsillitis, anemia, diarrhea, ear infection, infections, nasopharyngitis, hemoglobin decreased, and headache (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact Recordati Rare Diseases Inc. at 1-888-575-8344, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

- **Pregnancy:** No human data; decreased survival and growth in animal offspring. (8.1)
- **Nursing Mothers:** Breastfeeding is not recommended while taking CARBAGLU. (8.3)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 11/2017

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

- 1.1 Acute Hyperammonemia in Patients with NAGS Deficiency
- 1.2 Chronic Hyperammonemia in Patients with NAGS Deficiency

2 DOSAGE AND ADMINISTRATION

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

- 2.1 Recommended Dosage
- 2.2 Preparation and Administration

3 DOSAGE FORMS AND STRENGTHS

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

14 CLINICAL STUDIES

- 14.1 Responses of Patients with NAGS Deficiency to Acute and Chronic Treatment

6 ADVERSE REACTIONS

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

7 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Acute Hyperammonemia in Patients with NAGS Deficiency

CARBAGLU is indicated as an adjunctive therapy in pediatric and adult patients for the treatment of acute hyperammonemia due to the deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS). During acute hyperammonemic episodes, concomitant administration of CARBAGLU with other ammonia lowering therapies, such as alternate pathway medications, hemodialysis, and dietary protein restriction, is recommended.

1.2 Chronic Hyperammonemia in Patients with NAGS Deficiency

CARBAGLU is indicated as maintenance therapy in pediatric and adult patients for the treatment of chronic hyperammonemia due to deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS). During maintenance therapy, the concomitant use of other ammonia lowering therapies and protein restriction may be needed based on plasma ammonia levels.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

CARBAGLU should be initiated as soon as the diagnosis of NAGS deficiency is suspected, which may be as soon as at birth, and managed by a physician and medical team experienced in metabolic disorders.

Initial Dosage: The recommended initial daily dosage of CARBAGLU in pediatric and adult patients for acute hyperammonemia is 100 mg/kg to 250 mg/kg divided into 2 to 4 doses and rounded to the nearest 100 mg (i.e., half of a CARBAGLU tablet). Concomitant administration of other ammonia lowering therapies is recommended.

Maintenance Dosage: The recommended daily maintenance dosage of CARBAGLU in pediatric and adult patients is 10 mg/kg to 100 mg/kg divided into 2 to 4 doses and rounded to the nearest 100 mg (i.e., half of a CARBAGLU tablet).

Therapeutic Monitoring

Closely monitor plasma ammonia levels. Titrate the CARBAGLU dosage to maintain the plasma ammonia level within the normal range for the patient's age, taking into consideration their clinical condition (e.g., nutritional requirements, protein intake, growth parameters, etc.).

2.2 Preparation and Administration

- Disperse CARBAGLU tablets in water. Do not swallow whole or crushed.
- Mix each 200 mg tablet in a minimum of 2.5 mL of water to yield a concentration of 80 mg/mL.
- CARBAGLU tablets do not dissolve completely in water and undissolved particles of the tablet may remain in the mixing container.
- Take CARBAGLU immediately before meals or feedings.
- The CARBAGLU suspension has a slightly acidic taste.
- For all preparations, use in foods or liquids, other than water, has not been studied clinically and is not recommended.

Preparation for Oral Administration in Pediatric and Adult Patients

- Add about 2.5 mL of water into a small cup for each CARBAGLU tablet or each ½ CARBAGLU tablet needed for the prescribed dose.
- Add the CARBAGLU tablets to the water in the cup.
- Carefully stir the tablet and water mixture.
- Swallow the mixture immediately. Pieces of the tablet may remain in the cup.
- Rinse the cup with additional water and swallow the mixture immediately. Repeat as needed until no pieces of the tablet are left in the cup.

Preparation for Nasogastric Tube Administration in Pediatric and Adult Patients

For patients who have a nasogastric tube in place, CARBAGLU should be administered as follows:

- Add about 2.5 mL of water into a small cup for each CARBAGLU tablet or each ½ CARBAGLU tablet needed for the prescribed dose.
- Add the CARBAGLU tablets to the water in the cup.
- Carefully stir the tablet and water mixture.
- Draw up the mixture into a catheter-tip syringe.
- Administer the mixture immediately through the nasogastric (NG) tube. Pieces of the tablet may remain in the catheter-tip syringe or NG tube.

- Flush immediately with 1 to 2 mL of additional water to clear the NG tube.
- Flush the NG tube again, as needed, until no pieces of the tablet are left in the syringe or NG tube.

Preparation for Oral Administration Using an Oral Syringe in Pediatric Patients

For administration via oral syringe, CARBAGLU should be administered as follows:

- Add about 2.5 mL of water into a small cup for each CARBAGLU tablet or each ½ CARBAGLU tablet needed for the prescribed dose.
- Add the CARBAGLU tablets to the water in the cup.
- Carefully stir the tablet and water mixture.
- Draw up the mixture in an oral syringe and administer immediately. Pieces of the tablet may remain in the oral syringe.
- Refill the oral syringe with a minimum volume of water (1 to 2 mL) and administer immediately.
- Flush the oral syringe again, as needed, until no pieces of the tablet are left in the syringe.

3 DOSAGE FORMS AND STRENGTHS

CARBAGLU is a white and elongated 200 mg tablet for oral suspension, functionally scored and coded "C" on one side.

4 CONTRAINDICATIONS

None

5 WARNINGS AND PRECAUTIONS

5.1 Hyperammonemia

Any episode of acute symptomatic hyperammonemia should be treated as a life-threatening emergency. Treatment of severe hyperammonemia may require dialysis, preferably hemodialysis and/or hemofiltration, to reduce plasma ammonia concentration. Untreated hyperammonemia can result in brain damage and death, and prompt use of all therapies necessary to reduce plasma ammonia level is essential.

Since hyperammonemia in NAGS deficiency is the result of imbalance between ammonia detoxification capacity and protein catabolism, complete protein restriction during an acute hyperammonemic episode is recommended for no longer than 12 to 36 hours while maximizing caloric supplementation to reverse catabolism. Protein should be reintroduced as early as possible, following improvement of metabolic and clinical abnormalities in this setting. During long-term management, dietary protein restriction should be instituted to maintain blood ammonia level within an acceptable range for age.

Ongoing monitoring of plasma ammonia level, neurological status, growth parameters, protein intake/nutritional status (both during acute hyperammonemic episodes and long-term), and relevant laboratory tests in patients receiving CARBAGLU should be part of evaluating the clinical response to treatment.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Table 1 summarizes adverse reactions occurring in 2 or more patients treated with CARBAGLU in the retrospective case series (≥ 9%).

Table 1. Adverse Reactions Reported in ≥ 2 Patients in the Retrospective Case Series Treated with CARBAGLU

Adverse Reaction	Number of Patients (N)(%)
Vomiting	6 (26)
Abdominal pain	4 (17)
Pyrexia	4 (17)
Tonsillitis	4 (17)
Anemia	3 (13)
Diarrhea	3 (13)
Ear infection	3 (13)
Infections	3 (13)
Nasopharyngitis	3 (13)
Hemoglobin decreased	3 (13)
Headache	3 (13)
Dysguesia	2 (9)
Asthenia	2 (9)
Hyperhidrosis	2 (9)
Influenza	2 (9)
Pneumonia	2 (9)
Weight decreased	2 (9)
Anorexia	2 (9)
Somnolence	2 (9)
Rash	2 (9)

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C

There are no adequate and well controlled studies or available human data with CARBAGLU in pregnant women. Decreased survival and growth occurred in offspring born to animals that received arglumic acid at a dose approximately 38 times the maximum reported

human maintenance dose. Because untreated N-acetylglutamate synthase (NAGS) deficiency results in irreversible neurologic damage and death, women with NAGS must remain on treatment throughout pregnancy.

No effects on embryo-fetal development were observed in pregnant rats treated with up to 2000 mg/kg/day (approximately 38 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC [area under the plasma concentration-time curve]) from two weeks prior to mating through organogenesis or in pregnant rabbits treated with up to 1000 mg/kg/day (approximately 6 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC).

In a peri- and post-natal developmental study, female rats received oral arglumic acid from organogenesis through lactation at doses of 500 and 2000 mg/kg/day. Decreased growth of offspring was observed at 500 mg/kg/day and higher (approximately 38 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC) and reduction in offspring survival during lactation was observed at 2000 mg/kg/day (approximately 38 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC). No effects on physical and sexual development, learning and memory, or reproductive performance were observed through maturation of the surviving offspring at maternal doses up to 2000 mg/kg/day. The high dose (2000 mg/kg/day) produced maternal toxicity (impaired weight gain and approximately 10% mortality).

8.3 Nursing Mothers

It is not known whether CARBAGLU is excreted in human milk. Arglumic acid is excreted in rat milk, and an increase in mortality and impairment of body weight gain occurred in neonatal rats nursed by mothers receiving arglumic acid. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from CARBAGLU, breast-feeding is not recommended. Treatment is continuous and life-long for NAGS deficiency patients.

8.4 Pediatric Use

The efficacy of CARBAGLU for the treatment of hyperammonemia in patients with NAGS deficiency from birth to adulthood was evaluated in a retrospective review of the clinical course of 23 NAGS deficiency patients who all began CARBAGLU treatment during infancy or childhood. There are no apparent differences in clinical response between adults and pediatric NAGS deficiency patients treated with CARBAGLU. However, data are limited.

8.5 Geriatric Use

CARBAGLU has not been studied in the geriatric population. Therefore, the safety and effectiveness in geriatric patients have not been established.

10 OVERDOSAGE

One patient treated with 650 mg/kg/day of arglumic acid developed symptoms characterized as a monosodium glutamate intoxication-like syndrome: tachycardia, profuse sweating, increased bronchial secretion, increased body temperature and restlessness. These symptoms resolved upon reduction of dose.


Repeated oral dosing of arglumic acid at 2000 mg/kg/day was lethal to most neonatal rats within 2-3 days of treatment. The plasma concentrations that produced lethality were not measured. In adult rats, a single oral administration of arglumic acid was not lethal at doses up to 2800 mg/kg (approximately 20 times the maximum starting dose based on C_{max}).

11 DESCRIPTION

CARBAGLU tablets for oral suspension, contain 200 mg of arglumic acid. Arglumic acid, the active substance, is a Carbamoyl Phosphate Synthetase 1 (CPS 1) activator and is soluble in boiling water, slightly soluble in cold water, and practically insoluble in organic solvents.

Chemically arglumic acid is N-carbamoyl-L-glutamic acid or (2S)-2-(carbamoylamino) pentanedioic acid, with a molecular weight of 190.16.

The structural formula is:



Molecular Formula: $C_8H_{10}N_2O_5$

The inactive ingredients of CARBAGLU are croscarmellose sodium, hypromellose, microcrystalline cellulose, silica colloidal anhydrous, sodium lauryl sulfate, sodium stearyl fumarate.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Arglumic acid is a synthetic structural analogue of N-acetylglutamate (NAG) which is produced from glutamate and acetyl-CoA in a reaction catalyzed by N-acetylglutamate synthase (NAGS), a mitochondrial liver enzyme. NAG acts as an essential allosteric activator of Carbamoyl Phosphate Synthetase 1 (CPS 1), a mitochondrial liver enzyme which catalyzes the first reaction of the urea cycle. The urea cycle, whose role is the disposition of ammonia, includes a series of biochemical reactions in the liver resulting in the conversion of ammonia into urea, which is then excreted through the urine. Arglumic acid acts as a CPS1 activator in patients with NAGS deficiency, thereby removing the block in the urea cycle and facilitating ammonia detoxification and urea production.

12.2 Pharmacodynamics

In a retrospective review of the clinical course in 23 patients with NAGS deficiency, arglumic acid reduced plasma ammonia levels within 24 hours when administered with and without concomitant ammonia lowering therapies. No dose response relationship has been identified.

12.3 Pharmacokinetics

The pharmacokinetics of arglumic acid have been studied in healthy male subjects using both radiolabeled and non-radiolabeled arglumic acid.

Absorption

The median T_{max} of CARBAGLU was 3 hours (range: 2 to 4 hours). Absolute bioavailability has not been determined.

Distribution

The apparent volume of distribution was 2657 L (range: 1616-5797). Protein binding has not been determined.

Elimination

Metabolism

A proportion of carginic acid may be metabolized by the intestinal bacterial flora. The likely end product of carginic acid metabolism is carbon dioxide, eliminated through the lungs.

Excretion

Median value for the terminal half-life was 5.6 hours (range 4.3 to 9.5 hours), the apparent total clearance was 5.7 L/min (range 3.0 to 9.7 L/min), the renal clearance was 290 mL/min (range 204 to 445 mL/min), and the 24-hour urinary excretion was 4.5% of the dose (range 3.5 to 7.5%). Following administration of a single radiolabeled oral dose of 100 mg/kg of body weight, 9% of the dose was excreted unchanged in the urine and up to 60% of the dose was excreted unchanged in the feces.

Drug Interaction Studies

No drug interaction studies have been performed. Based on *in vitro* studies, CARBAGLU is not an inducer of CYP1A1/2, CYP2B6, CYP2C, and CYP3A4/5 enzymes, and not an inhibitor of CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1, and CYP3A4/5 enzymes.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

The carcinogenic potential of carginic acid was assessed in a 2-year carcinogenicity study in rats. Carginic acid was not tumorigenic at oral doses up to 1000 mg/kg/day (approximately 34 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC).

Carginic acid was negative in the Ames test, chromosomal aberration assay in human lymphocytes, and the *in vivo* micronucleus assay in rats.

There were no effects on fertility or reproductive performance in female rats at oral doses up to 2000 mg/kg/day (approximately 38 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC). In a separate study, mating and fertility were unaffected in male rats at oral doses up to 1000 mg/kg/day (approximately 34 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC).

14 CLINICAL STUDIES

14.1 Responses of Patients with NAGS Deficiency to Acute and Chronic Treatment

The efficacy of CARBAGLU in the treatment of hyperammonemia due to NAGS deficiency was evaluated in a retrospective review of the clinical course of 23 NAGS deficiency patients who received CARBAGLU treatment for a median of 7.9 years (range 0.6 to 20.8 years). Treatment with CARBAGLU was divided in two regimens. For acute treatment, patients received a total daily dose of 100 to 250 mg/kg per day primarily administered in 2 to 4 divided doses for the first few days of treatment. For maintenance treatment, the dosage was reduced over time based upon biochemical and clinical responses.

The demographics characteristics of the patient population are shown in Table 2.

Table 2. Baseline Characteristics of the 23 NAGS deficiency patients

Gender	Patients N=23	
	Male	14 (61%)
Age at initiation of CARBAGLU therapy (years)	Female	9 (39%)
	Mean (SD)	2 (4)
Age groups at initiation of CARBAGLU therapy	Min-Max	0-13
	< 30 days	9 (39%)
	>30 days - 11 months	9 (39%)
NAGS gene mutations by DNA testing	≥1 - 13 years	5 (22%)
	homozygous	14 (61%)
	heterozygous	4 (17%)
	Not available	5 (22%)
Patients current treatment status	On-going	18 (78%)
	Discontinued	5 (22%)

The clinical observations in the 23 patient case series were retrospective, unblinded and uncontrolled and preclude any meaningful formal statistical analyses of the data. However, short-term efficacy was evaluated using mean and median change in plasma ammonia levels from baseline to days 1 to 3. Persistence of efficacy was evaluated using long-term mean and median change in plasma ammonia level. Table 3 summarizes the plasma ammonia levels at baseline, days 1 to 3 post-CARBAGLU treatment, and long-term CARBAGLU treatment for 13 evaluable patients. Of the 23 NAGS deficiency patients who received treatment with CARBAGLU, a subset of 13 patients who had both well documented plasma ammonia levels prior to CARBAGLU treatment and after long-term treatment with CARBAGLU were selected for analysis.

All 13 patients had abnormal ammonia levels at baseline. The overall mean baseline plasma ammonia level was 271 micromol/L. By day 3, normal plasma ammonia levels were attained in patients for whom data were available. Long-term efficacy was measured using the last reported plasma ammonia level for each of the 13 patients analyzed (median length of treatment was 6 years; range 1 to 16 years). The mean and median ammonia levels were 23 micromol/L and 24 micromol/L, respectively, after a mean treatment duration of 8 years.


Table 3. Plasma ammonia levels at baseline and after treatment with CARBAGLU

Timepoint	Statistics (N = 13*)	Ammonia** (micromol/L)
Baseline (prior to first treatment with CARBAGLU)	N	13
	Mean (SD)	271 (359)
	Median	157
	Range	72-1428
	Missing Data	0
Day 1	N	10
	Mean (SD)	181 (358)
	Median	65
	Range	25-1190
	Missing Data	3
Day 2	N	8
	Mean (SD)	69 (78)
	Median	44
	Range	11-255
	Missing Data	5
Day 3	N	5
	Mean (SD)	27 (11)
	Median	25
	Range	12-42
	Missing Data	8
Long-term (last available value on CARBAGLU treatment)	N	13
	Mean (SD)	23 (7)
	Median	24
	Range	9-34
	Missing Data	0

* 13/23 patients with complete short-term and long-term plasma ammonia documentation
** Mean ammonia normal range: 5 to 50 micromol/L

The mean plasma ammonia level at baseline and the decline that is observed after treatment with CARBAGLU in 13 evaluable patients with NAGS deficiency is illustrated in Figure 1.

Figure 1. Ammonia response for 13 evaluable NAGS deficiency patients at baseline and after treatment with CARBAGLU



16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

CARBAGLU is a white and elongated 200 mg tablet for oral suspension, functionally scored and coded "C" on one side.

CARBAGLU is available in 5 or 60 tablets in a high density polyethylene bottle with child resistant polypropylene cap and desiccant unit.

NDC 52276-312-05 Bottles of 5 tablets

NDC 52276-312-60 Bottles of 60 tablets

Storage

Store in the original unopened container at 2 – 8°C (36 – 46°F).

After first opening of the container:

- Do not refrigerate, store at room temperature between 15 – 30°C (59 – 86°F).
- Keep the container tightly closed between openings in order to protect from moisture.
- Write the date of opening on the tablet container.
- Do not use after the expiration date stated on the tablet container.
- Discard one month after first opening.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Instructions for Use).

Preparation and Administration

- Disperse CARBAGLU tablets in water. Do not swallow whole or crushed.
- Take CARBAGLU immediately before meals or feedings.
- CARBAGLU tablets dispersed in water can be administered orally or via a nasogastric tube, as described in the *Instructions for Use*.

Storage

- Store UNOPENED container in a refrigerator at 2 to 8°C (36 to 46°F). After first opening of the container: do not refrigerate, store at room temperature between 15 to 30°C (59 to 86°F). Keep the container tightly closed in order to protect from moisture. Write the date of opening on the tablet container. Discard one month after first opening. Do not use after the expiration date stated on the tablet container.

Lactation

- Advise women not to breast-feed during treatment with CARBAGLU [see Use in Specific Populations (8.3)].

Supplied by:

Orphan Europe SARL
Puteaux, France



Licensed to and Distributed by:

Recordati Rare Diseases Inc.
Lebanon, NJ 08833



For drug or ordering information please call Accredo Health Group Inc., Customer Service at 1-888-454-8860.

Carbaglu
carglumic acid

Carbaglu® is a licensed trademark of Recordati Rare Diseases Inc.

This product label may have been updated. For the most recent prescribing information, please visit www.recordatirarediseases.com or www.carbaglu.net.

INSTRUCTIONS FOR USE CARBAGLU (CAR-buh-gloo) (carglumic acid) tablet for oral suspension

Important information:

- **CARBAGLU tablet for oral suspension (CARBAGLU tablet) must be mixed in water before taking.** CARBAGLU tablets should not be mixed in any other food or liquid.
- **Do not** swallow CARBAGLU tablets whole.
- **Do not** crush CARBAGLU tablets.
- Take CARBAGLU **right before** meals or feedings.
- The CARBAGLU tablet and water mixture has a slightly sour taste.

You may need to ask your healthcare provider or pharmacist for a medicine cup to measure the correct amount of water you will need to prepare your dose of CARBAGLU.

Taking CARBAGLU tablets by mouth using a cup:

Children and Adults

1. Add about 2.5 mL of water into a small cup for each CARBAGLU tablet, or each ½ CARBAGLU tablet, needed for the prescribed dose. For example, if the prescribed dose is 2 CARBAGLU tablets, add about 5 mL of water into the cup. If the prescribed dose is 2½ CARBAGLU tablets, add about 7.5 mL of water into the cup. Ask your healthcare provider if you are not sure of how much water you should use for the prescribed dose of CARBAGLU.
2. Place the prescribed number of CARBAGLU tablets into the water in the cup.
3. Carefully stir the CARBAGLU tablet and water mixture in the cup to avoid spilling the mixture. CARBAGLU tablets do not dissolve completely in water.
4. Swallow the CARBAGLU tablet and water mixture **right away**.
5. Pieces of the tablet may remain in the cup. Add more water to the cup to rinse the cup and swallow the mixture **right away**.
6. Repeat step 5 until there are no pieces of the tablet left in the cup.

Taking CARBAGLU tablets by mouth using an oral syringe:

Children

1. Add about 2.5 mL of water into a small cup for each CARBAGLU tablet, or each ½ CARBAGLU tablet, needed for the prescribed dose. For example, if the prescribed dose is 2 CARBAGLU tablets, add about 5 mL of water into the cup. If the prescribed dose is 2½ CARBAGLU tablets, add about 7.5 mL of water into the cup. Ask your healthcare provider if you are not sure of how much water you should use for the prescribed dose of CARBAGLU.
2. Place the prescribed number of CARBAGLU tablets into the water in the cup.
3. Carefully stir the CARBAGLU tablet and water mixture in the cup to avoid spilling the mixture. CARBAGLU tablets do not dissolve completely in water.
4. Draw up all of the CARBAGLU tablet and water mixture in the cup into an oral syringe.
5. Give your child the CARBAGLU tablet and water mixture **right away** by placing the tip of the oral syringe along the inner cheek of their mouth, on either the right or left side. Slowly push all the way down on the plunger to give the medicine.
6. Pieces of the tablet may remain in the oral syringe. Refill the oral syringe with at least 1 mL to 2 mL of water, and give your child the mixture **right away**.
7. Repeat step 6 until there are no pieces of the tablet left in the oral syringe.

Giving CARBAGLU tablets through a nasogastric (NG) tube:

Children and Adults

1. Add about 2.5 mL of water into a small cup for each CARBAGLU tablet, or each ½ CARBAGLU tablet, needed for the prescribed dose. For example, if the prescribed dose is 2 CARBAGLU tablets, add about 5 mL of water into the cup. If the prescribed dose is 2½ CARBAGLU tablets, add about 7.5 mL of water into the cup. Ask your healthcare provider if you are not sure of how much water you should use for the prescribed dose of CARBAGLU.
2. Place the prescribed number of CARBAGLU tablets into the water in the cup.
3. Carefully stir the CARBAGLU tablet and water mixture in the cup to avoid spilling the mixture. CARBAGLU tablets do not dissolve completely in water.
4. Draw up all of the CARBAGLU tablet and water mixture in the cup into a catheter-tip syringe.
5. Connect the catheter-tip syringe to the NG tube.
6. Give the CARBAGLU tablet and water mixture through the NG tube **right away**.
7. Pieces of the tablet may remain in the catheter-tip syringe or NG tube. Refill the catheter-tip syringe with 1 mL to 2 mL of water and flush the NG tube **right away**.
8. Repeat step 7 until there are no pieces of the tablet left in the catheter-tip syringe or NG tube.

How should I store CARBAGLU?

- **Before opening**, store CARBAGLU in a refrigerator between 36°F to 46°F (2°C to 8°C) in the container it comes in.
- **After opening**, store CARBAGLU at room temperature between 59°F to 86°F (15°C to 30°C). **Do not** store CARBAGLU in a refrigerator.
 - Keep CARBAGLU tablets in a tightly closed container to protect the tablets from moisture.
 - Write the date the CARBAGLU tablet container is opened on the container label. Throw away any unused tablets one month after opening the tablet container.
- Do not use CARBAGLU tablets after the expiration date on the tablet container.

Keep CARBAGLU and all medicines out of the reach of children.

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Distributed by:

Recordati Rare Diseases Inc.
Lebanon, NJ 08833



Issued: November 2017

Carbaglu
carglumic acid

Carbaglu® is a licensed trademark of Recordati Rare Diseases Inc.

PP-CBGL-US-0098