

Carbaglu[®]
(carglumic acid) tablets for oral suspension
200mg

Ordering CARBAGLU

for Acute and Chronic
Hyperammonemia
due to **NAGS Deficiency**



& Acute Hyperammonemia
due to **Propionic Acidemia**
or **Methylmalonic Acidemia**

INDICATIONS AND USAGE

CARBAGLU[®] (carglumic acid) tablets for oral suspension 200mg is a carbamoyl phosphate synthetase 1 (CPS1) activator indicated in pediatric and adult patients as:

- Adjunctive therapy to standard of care for the treatment of acute hyperammonemia due to N-acetylglutamate synthase (NAGS) deficiency.
- Maintenance therapy for the treatment of chronic hyperammonemia due to N-acetylglutamate synthase (NAGS) deficiency.
- Adjunctive therapy to standard of care for the treatment of acute hyperammonemia due to propionic acidemia (PA) or methylmalonic acidemia (MMA).

IMPORTANT SAFETY INFORMATION

- **NAGS deficiency:** Most common adverse reactions ($\geq 13\%$) are: vomiting, abdominal pain, pyrexia, tonsillitis, anemia, diarrhea, ear infection, infections, nasopharyngitis, hemoglobin decreased, and headache.
- **PA and MMA:** Most common adverse reactions ($\geq 5\%$) are neutropenia, anemia, vomiting, electrolyte imbalance, decreased appetite, hypoglycemia, lethargy/stupor, encephalopathy and pancreatitis/lipase increased.

Please see accompanying Full Prescribing Information, including Instructions for Use.

NAGS Deficiency: Treating Hyperammonemia with CARBAGLU®

Treatment Guidelines

- Initiate CARBAGLU as soon as the diagnosis of NAGS deficiency is suspected, which may be as soon as at birth.
- CARBAGLU treatment should be managed by a physician and medical team experienced in metabolic disorders.
- During acute hyperammonemic episodes, administer CARBAGLU with other ammonia lowering therapies, such as alternate pathway medications, hemodialysis, and protein restriction.
- During maintenance therapy, the concomitant use of other ammonia lowering therapies and protein restriction may be needed based on plasma ammonia levels. Closely monitor plasma ammonia levels. Titrate the CARBAGLU dosage to maintain the plasma ammonia level within the normal range for the patient's age, taking into consideration their clinical condition (e.g., nutritional requirements, protein intake, growth parameters, etc.).

Preparation and Administration

- Disperse CARBAGLU tablets in water. Do not swallow whole or crushed. Mix each whole or partial 200 mg tablet in a minimum of 2.5 mL of water.
- CARBAGLU can be taken orally via a cup or oral syringe or via a nasogastric or gastrostomy tube.

IMPORTANT SAFETY INFORMATION

- Most common adverse reactions ($\geq 13\%$) are: vomiting, abdominal pain, pyrexia, tonsillitis, anemia, diarrhea, ear infection, infections, nasopharyngitis, hemoglobin decreased, and headache.
- **To report SUSPECTED ADVERSE REACTIONS, contact Recordati Rare Diseases Inc. at 1-888-575-8344, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.**
- If CARBAGLU is administered during pregnancy to women with NAGS deficiency, health care providers should report CARBAGLU exposure to the pregnancy pharmacovigilance program by calling 1-888-575-8344.

Please see accompanying Full Prescribing Information, including Instructions for Use.

Carbaglu®
(carglumic acid) tablets for oral suspension
200mg

Dosing CARBAGLU: NAGS Deficiency

NAGS Deficiency Dosing and Administration*

Patient Type	Recommended Dose	Frequency & Timing	Dose Titration - Yes/No
Acute Hyperammonemia Pediatric & Adult Patients	100 mg/kg/day to 250 mg/kg/day	Divide total daily dose into 2 to 4 doses to be given immediately before meals or feedings. Round dose to the nearest 100 mg (i.e., half of a CARBAGLU tablet).	No.
Maintenance for Chronic Hyperammonemia Pediatric & Adult Patients	10 mg/kg/day to 100 mg/kg/day	Divide total daily dose into 2 to 4 doses to be given immediately before meals or feedings. Round dose to the nearest 100 mg (i.e., half of a CARBAGLU tablet).	Yes. Titrate to maintain the plasma ammonia level within the normal range for the patient's age, taking into consideration their clinical condition.

*See the next page for dosing adjustments for NAGS deficiency patients with renal impairment.

NAGS Deficiency Sample Dosing Guides to Help with Emergency Ordering

Patient – Weight	Acute initial dose (min/max)	Total daily mg	# of 200-mg tablets per day	3-day supply for acute hyperammonemic episode # of 60-tablet and/or 5-tablet bottles
For example: Neonate 4 kg (8.82 lbs)	100 mg/kg	400 mg	2 tablets	2 bottles of 5 tablets
	250 mg/kg	1000 mg	5 tablets	3 bottles of 5 tablets
For example: 4-year-old 16 kg (35.27 lbs)	100 mg/kg	1600 mg	8 tablets	5 bottles of 5 tablets
	250 mg/kg	4000 mg	20 tablets	1 bottle of 60 tablets
For example: Adult 68 kg (149.91 lbs)	100 mg/kg	6800 mg	34 tablets	1 bottle of 60 tablets and 9 bottles of 5 tablets
	250 mg/kg	17000 mg	85 tablets	4 bottles of 60 tablets and 3 bottles of 5 tablets

IMPORTANT SAFETY INFORMATION

- If CARBAGLU is administered during pregnancy to women with NAGS deficiency, health care providers should report CARBAGLU exposure to the pregnancy pharmacovigilance program by calling 1-888-575-8344.

Please see accompanying Full Prescribing Information, including Instructions for Use.

Carbaglu[®]
(carglumic acid) tablets for oral suspension
200mg

Dosing CARBAGLU in NAGS Deficiency: Adjustments for Renal Impairment

Mild Renal Impairment (eGFR 60-89 mL/min/1.73 m²)

Patient Type	Recommended Dose
All	No dosage adjustment is warranted. Use guides on page 2.

Moderate Renal Impairment (eGFR 30-59 mL/min/1.73 m²)

Patient Type	Recommended Dose	Frequency & Timing
Acute Hyperammonemia due to NAGS Deficiency	50 mg/kg/day to 125 mg/kg/day	Divide the daily dosage into 2 to 4 doses. Round dose to the nearest 50 mg (i.e., one-quarter of a CARBAGLU tablet).
Chronic Hyperammonemia due to NAGS Deficiency	5 mg/kg/day to 50 mg/kg/day	Divide the daily dosage into 2 to 4 doses. Round dose to the nearest 50 mg (i.e., one-quarter of a CARBAGLU tablet).

Severe Renal Impairment (eGFR ≤29 mL/min/1.73 m²)

Patient Type	Recommended Dose	Frequency & Timing
Acute Hyperammonemia due to NAGS Deficiency	15 mg/kg/day to 60 mg/kg/day	Divide the daily dosage into 2 to 4 doses. Round dose to the nearest 50 mg (i.e., one-quarter of a CARBAGLU tablet).
Chronic Hyperammonemia due to NAGS Deficiency	2 mg/kg/day to 25 mg/kg/day	Divide the daily dose into 2 to 4 doses. Round to the nearest 50 mg (i.e., one-quarter of a CARBAGLU tablet).

IMPORTANT SAFETY INFORMATION

- Contraindications: None.
- Most common adverse reactions (≥13%) are: vomiting, abdominal pain, pyrexia, tonsillitis, anemia, diarrhea, ear infection, infections, nasopharyngitis, hemoglobin decreased, and headache.

Please see accompanying Full Prescribing Information, including Instructions for Use.

Carbaglu[®]
(carglumic acid) tablets for oral suspension
200mg

PA or MMA: Treating Acute Hyperammonemia with CARBAGLU®

Treatment Guidelines

- Initiate CARBAGLU for the treatment of acute hyperammonemia in patients with a suspected or confirmed diagnosis of propionic acidemia (PA) or methylmalonic acidemia (MMA).
- During acute hyperammonemic episodes, administer CARBAGLU with other ammonia lowering therapies, such as intravenous glucose, insulin, L-carnitine, protein restriction, and dialysis.

Preparation and Administration

- Disperse CARBAGLU tablets in water. Do not swallow whole or crushed. Mix each whole or partial 200 mg tablet in a minimum of 2.5 mL of water.
- CARBAGLU can be taken orally via a cup or oral syringe or via a nasogastric or gastrostomy tube.

IMPORTANT SAFETY INFORMATION

- Most common adverse reactions ($\geq 5\%$) are neutropenia, anemia, vomiting, electrolyte imbalance, decreased appetite, hypoglycemia, lethargy/stupor, encephalopathy and pancreatitis/lipase increased.
- To report **SUSPECTED ADVERSE REACTIONS**, contact **Recordati Rare Diseases Inc. at 1-888-575-8344, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.**

Please see accompanying Full Prescribing Information, including Instructions for Use.

Carbaglu®
(carglumic acid) tablets for oral suspension
200mg

Dosing CARBAGLU: PA or MMA

PA or MMA Dosing and Administration*

Patient Type	Recommended Dose	Frequency & Timing	Duration
Acute Hyperammonemia Pediatric & Adult Patients	150 mg/kg/day for patients ≤15 kg 3.3 g/m ² /day for patients >15 kg	Divide the daily dosage into 2 equal doses. Administer doses 12 hours apart. Round up dose to the nearest 50 mg (i.e., one quarter of a CARBAGLU tablet).	Continue CARBAGLU treatment until the patient's ammonia level is <50 micromol/L and for a maximum duration of 7 days.

*See the next page for dosing adjustments for PA or MMA patients with renal impairment.

PA or MMA Sample Dosing Guides to Help with Emergency Ordering

Patient – Weight or body surface area	Acute dose	Total daily mg	# of 200-mg tablets per day	3-day supply for acute hyperammonemic episode # of 60-tablet and/or 5-tablet bottles
For example: Neonate 4 kg (8.82 lbs)	150 mg/kg/day	600 mg	3 tablets	2 bottles of 5 tablets
For example: 4-year-old 16 kg (35.27 lbs) 102 cm (40.16 in)	3.3 g/m ² /day	2221.89 mg [†]	11.5 tablets	7 bottles of 5 tablets
For example: Adult 68 kg (149.91 lbs) 175 cm (68.90 in)	3.3 g/m ² /day	6000 mg*	30 tablets	1 bottle of 60 tablets and 6 bottles of 5 tablets

*Formula used: Mosteller body surface area (BSA). Online BSA calculators are available. Correlation between formulas is generally high.

[†]2300 mg when divided into two doses and rounded up to the nearest 50 mg.

IMPORTANT SAFETY INFORMATION

- Contraindications: None.

Carbaglu[®]
 (carglumic acid) tablets for oral suspension
 200mg

Please see accompanying Full Prescribing Information, including Instructions for Use.

Dosing CARBAGLU in PA or MMA: Adjustments for Renal Impairment

Mild Renal Impairment (eGFR 60-89 mL/min/1.73 m²)

Patient Type	Recommended Dose
All	No dosage adjustment is warranted. Use guides on page 5.

Moderate Renal Impairment (eGFR 30-59 mL/min/1.73 m²)

Patient Type	Recommended Dose	Frequency & Timing
Acute Hyperammonemia due to PA or MMA	<ul style="list-style-type: none"> 75 mg/kg/day for patients weighing ≤ 15 kg 1.7 g/m²/day for patients weighing > 15 kg 	Divide the daily dosage into 2 equal doses. Round up dose to the nearest 50 mg (i.e., one-quarter of a CARBAGLU tablet). Administer each dose 12 hours apart.

Severe Renal Impairment (eGFR ≤29 mL/min/1.73 m²)

Patient Type	Recommended Dose	Frequency & Timing
Acute Hyperammonemia due to PA or MMA	<ul style="list-style-type: none"> 25 mg/kg/day for patients weighing ≤ 15 kg 0.55 g/m²/day for patients weighing > 15 kg 	Divide the daily dosage into 2 equal doses. Round up dose to the nearest 50 mg (i.e., one-quarter of a CARBAGLU tablet). Administer each dose 12 hours apart.

IMPORTANT SAFETY INFORMATION

- Contraindications: None.
- Most common adverse reactions (≥5%) are neutropenia, anemia, vomiting, electrolyte imbalance, decreased appetite, hypoglycemia, lethargy/stupor, encephalopathy and pancreatitis/lipase increased.

Please see accompanying Full Prescribing Information, including Instructions for Use.

Carbaglu[®]
(carglumic acid) tablets for oral suspension
200mg

Ordering CARBAGLU® (carglumic acid) tablets for oral suspension 200mg

Exclusive Distribution Through Accredo

Representatives at Accredo Health Group, Inc. are committed to helping you and your patients through the ordering process. CARBAGLU is not available in retail pharmacies.

Patient Home Delivery

1. Download the CARBAGLU Prescription & Enrollment Form from the Accredo website (www.accredo.com).
2. Complete the CARBAGLU Prescription & Enrollment Form and fax it to the Accredo specialty pharmacy department at 1-888-454-8488.
3. A CARBAGLU specialty pharmacy representative may call to verify information and determine next steps. In some cases, prior authorization may be necessary.
4. Questions? Contact a CARBAGLU specialty pharmacy representative at 1-888-454-8860.

Hospital Orders – Emergency or Inpatient

1. Alert your hospital pharmacy that your patient requires CARBAGLU. Send an order for CARBAGLU to your hospital pharmacy and specify whether it is a STAT order (required in 6 hours or less*).
2. Ask your hospital pharmacy to call the wholesale department at Accredo at 1-877-900-9223 to place the order.
3. The wholesale department at Accredo will work with your hospital pharmacy to obtain payment information, establish shipping timelines, and verify the delivery address.
4. The Accredo specialty pharmacy team will follow up in 1-2 days to help set up patient home delivery of CARBAGLU, if applicable.

*Delivery time frame is weather dependent and is not guaranteed.

Keep a 5-count bottle of CARBAGLU in stock to allow for immediate treatment.

Return policy: All new orders of 5-count unopened bottles of CARBAGLU can be returned for credit any time prior to the expiration date or up to 12 months past the expiration date.†

†Not applicable for CARBAGLU 60-count bottles. Product shipped as an emergency or STAT order is not eligible for credit unless damaged in transit. This policy covers products purchased after July 1, 2019. Unopened bottles have up to a 3-year expiration date and must be refrigerated. Other terms and conditions may apply.

Questions? Contact a CARBAGLU specialty pharmacy representative at 1-888-454-8860.

Financial Assistance Programs

For patients experiencing financial hardship, Recordati Rare Diseases Inc. supports a Patient Assistance Program and a Co-Pay Assistance Program, administered by Accredo. For more information, call: 1-888-454-8860.

FOR MORE INFORMATION ABOUT CARBAGLU, CALL 1-888-454-8860

**Be prepared for patients who may need CARBAGLU in emergency situations.
Keep a 5-count bottle of CARBAGLU in stock to allow for immediate treatment.**



Unopened 5-count bottles of CARBAGLU can be returned for credit any time prior to the expiration date or up to 12 months past the expiration date.*

*Not applicable for CARBAGLU 60-count bottles. Product shipped as an emergency or STAT order is not eligible for credit unless damaged in transit. This policy covers products purchased after July 1, 2019. Unopened bottles have up to a 3-year expiration date and must be refrigerated. Other terms and conditions may apply.

**Please see accompanying Full Prescribing Information,
including Instructions for Use.**

For more information, visit www.CARBAGLU.com

CARBAGLU is a licensed trademark of Recordati Rare Diseases Inc. Other trademarks, registered or otherwise, are the property of their respective owner(s).

© 2021 Recordati Rare Diseases Inc.

Recordati Rare Diseases Inc. • Lebanon, NJ 08833

www.recordatirarediseases.com/us PP-CBGL-US-0244v2

 **RECORDATI
RARE DISEASES**
GROUP
Focused on the Few®

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use CARBAGLU safely and effectively. See full prescribing information for CARBAGLU.

CARBAGLU[®] (carglumic acid) tablets for oral suspension

Initial U.S. Approval: 2010

RECENT MAJOR CHANGES

Indications and Usage (1.2)	1/2021
Dosage and Administration (2.2, 2.4)	1/2021
Dosage and Administration, Renal Impairment (2.3)	8/2021

INDICATIONS AND USAGE

CARBAGLU is a carbamoyl phosphate synthetase 1 (CPS 1) activator indicated in pediatric and adult patients as:

- Adjunctive therapy to standard of care for the treatment of acute hyperammonemia due to N-acetylglutamate synthase (NAGS) deficiency. (1.1)
- Maintenance therapy for the treatment of chronic hyperammonemia due to NAGS deficiency. (1.1)
- Adjunctive therapy to standard of care for the treatment of acute hyperammonemia due to propionic acidemia (PA) or methylmalonic acidemia (MMA). (1.2)

DOSAGE AND ADMINISTRATION

Acute Hyperammonemia due to NAGS deficiency (2.1)

- The recommended pediatric and adult dosage is 100 mg/kg/day to 250 mg/kg/day divided into 2 to 4 doses and rounded to the nearest 100 mg.

Chronic Hyperammonemia due to NAGS deficiency (2.1)

- The recommended pediatric and adult dosage is 10 mg/kg/day to 100 mg/kg/day divided into 2 to 4 doses and rounded to the nearest 100 mg.

Therapeutic Monitoring for NAGS Deficiency (2.1)

- Closely monitor plasma ammonia and titrate dosage to maintain the ammonia level within normal range for the patient's age, taking into consideration their clinical condition.

Acute Hyperammonemia due to PA or MMA (2.2)

- The recommended pediatric and adult dosage is:
 - 150 mg/kg/day for patients less than or equal to 15 kg
 - 3.3 g/m²/day for patients greater than 15 kg
- Divide the daily dosage into two equal doses and round up to the nearest 50 mg; administer each dose 12 hours apart.
- Continue treatment until ammonia level is less than 50 micromol/L and for a maximum duration of 7 days.

See Full Prescribing Information for Instructions on Dosage Adjustment in Patients with Renal Impairment (2.3)

Preparation and Administration (2.4)

- Disperse CARBAGLU tablets in water. Do not swallow whole or crushed.
- Take immediately before meals or feedings.
- For additional instructions on preparation and administration orally or through a nasogastric tube or gastrostomy tube, see Full Prescribing Information.

DOSAGE FORMS AND STRENGTHS

Tablets for oral suspension: 200 mg, functionally scored. (3)

CONTRAINDICATIONS

None. (4)

ADVERSE REACTIONS

- **NAGS deficiency:** Most common adverse reactions (≥13%) are vomiting, abdominal pain, pyrexia, tonsillitis, anemia, diarrhea, ear infection, infections, nasopharyngitis, hemoglobin decreased, and headache. (6.1)
- **PA and MMA:** Most common adverse reactions (≥5%) are neutropenia, anemia, vomiting, electrolyte imbalance, decreased appetite, hypoglycemia, lethargy/stupor, encephalopathy and pancreatitis/lipase increased. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Recordati Rare Diseases Inc. at 1-888-575-8344, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 8/2021

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 1.1 Acute and Chronic Hyperammonemia due to N-acetylglutamate Synthase (NAGS) Deficiency
- 1.2 Acute Hyperammonemia due to Propionic Acidemia (PA) or Methylmalonic Acidemia (MMA)

2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Dosage for Acute or Chronic Hyperammonemia due to NAGS Deficiency
- 2.2 Recommended Dosage for Acute Hyperammonemia due to PA or MMA
- 2.3 Dosage Adjustment in Patients with Renal Impairment
- 2.4 Preparation and Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy

- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Acute and Chronic Hyperammonemia due to NAGS Deficiency
- 14.2 Acute Hyperammonemia due to PA and MMA

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Acute and Chronic Hyperammonemia due to N-acetylglutamate Synthase (NAGS) Deficiency

CARBAGLU is indicated in pediatric and adult patients as:

- Adjunctive therapy to standard of care for the treatment of acute hyperammonemia due to NAGS deficiency.
- Maintenance therapy for the treatment of chronic hyperammonemia due to NAGS deficiency.

1.2 Acute Hyperammonemia due to Propionic Acidemia (PA) or Methylmalonic Acidemia (MMA)

CARBAGLU is indicated in pediatric and adult patients as adjunctive therapy to standard of care for the treatment of acute hyperammonemia due to PA or MMA.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage for Acute or Chronic Hyperammonemia due to NAGS Deficiency

Treatment Initiation

Initiate CARBAGLU treatment as soon as the diagnosis of NAGS deficiency is suspected, which may be as soon as at birth, and managed by a physician and medical team experienced in metabolic disorders.

Dosage for Acute Hyperammonemia due to NAGS Deficiency

- The recommended daily dosage of CARBAGLU in pediatric and adult patients for acute hyperammonemia due to NAGS deficiency is 100 mg/kg to 250 mg/kg.
- Divide the daily dosage into 2 to 4 doses and round to the nearest 100 mg (i.e., half of a CARBAGLU tablet).
- During acute hyperammonemic episodes, administer CARBAGLU with other ammonia lowering therapies, such as alternate pathway medications, hemodialysis, and protein restriction.

Dosage for Chronic Hyperammonemia due to NAGS Deficiency

- The recommended daily dosage of CARBAGLU in pediatric and adult patients for chronic hyperammonemia due to NAGS deficiency is 10 mg/kg to 100 mg/kg.
- Divide the daily dosage into 2 to 4 doses and round to the nearest 100 mg (i.e., half of a CARBAGLU tablet).
- During maintenance therapy, the concomitant use of other ammonia lowering therapies and protein restriction may be needed based on plasma ammonia levels.

Therapeutic Monitoring

Closely monitor plasma ammonia levels. Titrate the CARBAGLU dosage to maintain the plasma ammonia level within the normal range for the patient's age, taking into consideration their clinical condition (e.g., nutritional requirements, protein intake, growth parameters, etc.).

Adjust the recommended dosage in patients with moderate or severe renal impairment [see Dosage and Administration (2.3)].

2.2 Recommended Dosage for Acute Hyperammonemia due to PA or MMA

Treatment Initiation

Initiate CARBAGLU for the treatment of acute hyperammonemia in patients with a suspected or confirmed diagnosis of PA or MMA.

Dosage for Acute Hyperammonemia due to PA or MMA

- The recommended daily dosage of CARBAGLU in pediatric and adult patients with acute hyperammonemia due to PA or MMA is:
 - 150 mg/kg/day for patients less than or equal to 15 kg
 - 3.3 g/m²/day for patients greater than 15 kg
- Divide the daily dosage into 2 equal doses and round up to the nearest 50 mg (i.e., one-quarter of a CARBAGLU tablet). Administer doses 12 hours apart.
- Continue CARBAGLU treatment until the patient's ammonia level is less than 50 micromol/L and for a maximum duration of 7 days.
- During acute hyperammonemic episodes, administer CARBAGLU with other ammonia lowering therapies, such as intravenous glucose, insulin, L-carnitine, protein restriction, and dialysis.

Adjust the recommended dosage in patients with moderate or severe renal impairment [see Dosage and Administration (2.3)].

2.3 Dosage Adjustment in Patients with Renal Impairment

No dosage adjustment is warranted in patients with mild renal impairment (eGFR 60-89 mL/min/1.73 m²). The recommended dosage of CARBAGLU in patients with moderate or severe renal impairment is shown below.

	Moderate Renal Impairment (eGFR 30-59 mL/min/1.73 m ²)	Severe Renal Impairment (eGFR ≤29 mL/min/1.73 m ²)
Acute Hyperammonemia due to NAGS Deficiency	50 mg/kg/day to 125 mg/kg/day divided into 2 to 4 doses and rounded to the nearest 50 mg (i.e., one-quarter of a CARBAGLU tablet)	15 mg/kg/day to 60 mg/kg/day divided into 2 to 4 doses and rounded to the nearest 50 mg (i.e., one-quarter of a CARBAGLU tablet)
Chronic Hyperammonemia due to NAGS Deficiency	5 mg/kg/day to 50 mg/kg/day divided into 2 to 4 doses and rounded to the nearest 50 mg (i.e., one-quarter of a CARBAGLU tablet)	2 mg/kg/day to 25 mg/kg/day divided into 2 to 4 doses and rounded to the nearest 50 mg (i.e., one-quarter of a CARBAGLU tablet)
Acute Hyperammonemia due to PA or MMA	<ul style="list-style-type: none">◦ 75 mg/kg/day for patients weighing ≤ 15 kg◦ 1.7 g/m²/day for patients weighing > 15 kg Divide daily dosage into 2 equal doses and round up to the nearest 50 mg (i.e., one-quarter of a CARBAGLU tablet); administer each dose 12 hours apart.	<ul style="list-style-type: none">◦ 25 mg/kg/day for patients weighing ≤ 15 kg◦ 0.55 g/m²/day for patients weighing > 15 kg Divide daily dosage into 2 equal doses and round up to the nearest 50 mg (i.e., one-quarter of a CARBAGLU tablet); administer each dose 12 hours apart.

2.4 Preparation and Administration

Overview

- Disperse CARBAGLU tablets in water. Do not swallow whole or crushed.
- CARBAGLU tablets do not dissolve completely in water, and undissolved particles of the tablet may remain in the mixing container.
- Take CARBAGLU immediately before meals or feedings.
- The CARBAGLU suspension has a slightly acidic taste.
- For all preparations, use in foods or liquids other than water has not been studied and is not recommended.

Oral Administration

For oral administration, administer CARBAGLU as follows:

- Add a minimum of 2.5 mL of water into a small cup for each CARBAGLU tablet or each ½ or ¼ CARBAGLU tablet needed for the prescribed dose.
- Add the CARBAGLU tablets to the water in the cup.
- Carefully stir the tablet and water mixture.

- Swallow the mixture immediately. Pieces of the tablet may remain in the cup.
- Rinse the cup with additional water and swallow the mixture immediately. Repeat as needed until no pieces of the tablet are left in the cup.

Use of an Oral Syringe for Oral Administration

For administration via an oral syringe, administer CARBAGLU as follows:

- Add a minimum of 2.5 mL of water into a small cup for each CARBAGLU tablet or each ½ or ¼ CARBAGLU tablet needed for the prescribed dose.
- Add the CARBAGLU tablets to the water in the cup.
- Carefully stir the tablet and water mixture.
- Draw up the mixture in an oral syringe and administer immediately. Pieces of the tablet may remain in the oral syringe.
- Refill the oral syringe with a minimum volume of water (1 mL to 2 mL) and administer immediately.
- Flush the oral syringe again, as needed, until no pieces of the tablet are left in the syringe.

Use of Nasogastric Tube (NG Tube) or Gastrostomy Tube (G-Tube) for Feeding Tube Administration

For patients who have a NG tube or G-tube in place, administer CARBAGLU as follows:

- Add a minimum of 2.5 mL of water into a small cup for each CARBAGLU tablet or each ½ or ¼ CARBAGLU tablet needed for the prescribed dose.
- Add the CARBAGLU tablets to the water in the cup.
- Carefully stir the tablet and water mixture.
- Draw up the mixture into a catheter-tip syringe.
- Administer the mixture immediately through the NG tube or G-tube. Pieces of the tablet may remain in the catheter-tip syringe or the feeding tube.
- Flush immediately with 1 to 2 mL of additional water to clear the NG tube or G-tube.
- Flush the NG tube or G-tube again, as needed, until no pieces of the tablet are left in the syringe or the feeding tube.

3 DOSAGE FORMS AND STRENGTHS

CARBAGLU is a white and elongated 200 mg tablet for oral suspension, functionally scored with 3 lines for splitting into 4 equal portions, and coded "C" on one side.

4 CONTRAINDICATIONS

None.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Acute and Chronic Hyperammonemia due to NAGS Deficiency

In a retrospective case series of 23 NAGS deficiency patients treated with CARBAGLU, 17 of the 23 patients reported any adverse reaction. The most common adverse reactions (occurring in ≥ 13% of patients) were vomiting, abdominal pain, pyrexia, tonsillitis, anemia, diarrhea, ear infection, infections, nasopharyngitis, hemoglobin decreased, and headache.

Table 1 summarizes adverse reactions occurring in 2 or more patients with NAGS deficiency treated with CARBAGLU in the retrospective case series (≥ 9%).

Table 1: Adverse Reactions Reported in ≥ 2 Patients with NAGS deficiency Treated with CARBAGLU in the Retrospective Case Series

Adverse Reaction	Number of Patients (N) (%)
Vomiting	6 (26)
Abdominal pain	4 (17)
Pyrexia	4 (17)
Tonsillitis	4 (17)
Anemia	3 (13)
Diarrhea	3 (13)
Ear infection	3 (13)
Infections	3 (13)
Nasopharyngitis	3 (13)
Hemoglobin decreased	3 (13)
Headache	3 (13)
Dysgeusia	2 (9)
Asthenia	2 (9)
Hyperhidrosis	2 (9)
Influenza	2 (9)
Pneumonia	2 (9)
Weight decreased	2 (9)
Anorexia	2 (9)
Somnolence	2 (9)
Rash	2 (9)

Acute Hyperammonemia due to PA and MMA

In a randomized, double-blind, placebo-controlled clinical trial, 24 patients (15 with PA and 9 with MMA) experienced a total of 90 hyperammonemic episodes which were randomized 1:1 to be treated with either CARBAGLU or placebo, each in addition to standard-of-care therapy, with randomization based on each hyperammonemic episode. The average patient age (SD) was 9.2 years (7.7) and 12 (50 %) were males.

CARBAGLU was administered at a dosage of 150 mg/kg/day for patients ≤ 15 kg or 3.3 g/m²/day for patients > 15 kg, divided into 2 doses, for a median duration of 4.14 days.

At least 1 adverse reaction was reported during the course of hyperammonemic episodes in 42.2% of hyperammonemic episodes. The most common adverse reactions ($\geq 5\%$) during hyperammonemic episodes were neutropenia, anemia, vomiting, electrolyte imbalance, decreased appetite, hypoglycemia, lethargy/stupor, encephalopathy and pancreatitis/lipase increased.

Table 2 summarizes adverse reactions ($\geq 2\%$) during hyperammonemic episodes in patients with PA or MMA treated with CARBAGLU or placebo.

Table 2: Adverse Reactions ($\geq 2\%$) During Hyperammonemic Episodes in Patients with PA or MMA Treated with CARBAGLU or Placebo

Adverse Reaction	Treatment Arm	
	CARBAGLU N=42 episodes	PLACEBO N=48 episodes
	N (%)	N (%)
Neutropenia	6 (14)	4 (8)
Anemia	5 (12)	4 (8)
Vomiting	3 (7)	1 (2)
Electrolyte imbalance	3 (7)	2 (4)
Decreased appetite	2 (5)	1 (2)
Hypoglycemia	2 (5)	1 (2)
Lethargy/Stupor	2 (5)	1 (2)
Encephalopathy	2 (5)	0 (0)
Pancreatitis/Lipase increased	2 (5)	0 (0)
Cardiomyopathy	1 (2)	0 (0)
Alanine aminotransferase increased	1 (2)	0 (0)
Aspartate aminotransferase increased	1 (2)	0 (0)
Infusion site extravasation	1 (2)	0 (0)
White blood cell count increased	1 (2)	0 (0)
Behavior disorder	1 (2)	0 (0)
Sleep disorder	1 (2)	0 (0)
Apnea	1 (2)	0 (0)
Hyperventilation	1 (2)	0 (0)

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of CARBAGLU. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or to establish a causal relationship to drug exposure.

Psychiatric disorders: mania

Skin and subcutaneous tissue disorders: pruritus, rash including rash erythematous, rash maculopapular, rash pustular

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women with NAGS deficiency exposed to CARBAGLU. If CARBAGLU is administered during pregnancy, health care providers should report CARBAGLU exposure by calling 1-888-575-8344.

Risk Summary

Although rare case reports of CARBAGLU use in pregnant women are insufficient to inform a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes, untreated NAGS deficiency, PA and MMA can result in irreversible neurologic damage and death in pregnant women [see *Clinical Considerations*].

In an animal reproduction study, decreased survival and growth occurred in offspring born to rats that received carglumic acid at a dose approximately 38 times the maximum reported human maintenance dose.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, miscarriage, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Clinical Considerations

Disease-associated maternal and/or embryo/fetal risk

Pregnant women with urea cycle disorders, PA, and MMA may experience an increase in catabolic stress which can trigger a hyperammonemic crisis both in the intrapartum and in the post-partum (3-14 days post-partum) periods. Maternal complications related to hyperammonemic crisis can include neurological impairment, coma and in some cases death.

Data

Animal Data

No effects on embryo-fetal development were observed in pregnant rats treated with up to 2000 mg/kg/day (approximately 38 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC [area under the plasma concentration-time curve]) from two weeks prior to mating through organogenesis or in pregnant rabbits treated with up to 1000 mg/kg/day (approximately 6 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC) during organogenesis.

In a pre- and post-natal developmental study, female rats received oral carglumic acid from organogenesis through lactation at doses of 500 mg/kg/day and 2000 mg/kg/day. Decreased growth of offspring was observed at 500 mg/kg/day and higher (approximately 38 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC), and reduction in offspring survival during lactation was observed at 2000 mg/kg/day (approximately 38 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC). No effects on physical and sexual development, learning and memory, or reproductive performance were observed through maturation of the surviving offspring at maternal doses up to 2000 mg/kg/day. The high dose (2000 mg/kg/day) produced maternal toxicity (impaired weight gain and approximately 10% mortality).

8.2 Lactation

Risk Summary

It is not known whether carglumic acid is present in human milk. There are no available data on the effects of carglumic acid on the breastfed infant or the effects on milk production. Carglumic acid is present in milk from treated rats. When a drug is present in animal milk, it is likely that the drug will be present in human milk.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for CARBAGLU and any potential adverse effects on the breastfed child from CARBAGLU or from the underlying maternal condition.

8.4 Pediatric Use

The safety and effectiveness of CARBAGLU for the treatment of pediatric patients (birth to 17 years of age) with acute or chronic hyperammonemia due to NAGS deficiency and acute hyperammonemia due to PA or MMA have been established, and the information on these uses are discussed throughout the labeling. There are insufficient data to determine if there is a difference in clinical or biochemical responses between adult and pediatric patients treated with CARBAGLU.

8.5 Geriatric Use

Clinical studies of CARBAGLU did not include patients 65 years of age and older.

8.6 Renal Impairment

Plasma concentrations of carglumic acid increased in patients with renal impairment [see *Clinical Pharmacology* (12.3)]. Reduce the CARBAGLU dosage in patients with moderate or severe renal impairment [see *Dosage and Administration* (2.3)]. The pharmacokinetics of carglumic acid have not been evaluated in patients with end stage renal disease.

10 OVERDOSAGE


One patient treated with 650 mg/kg/day of CARBAGLU developed symptoms resembling monosodium glutamate intoxication-like syndrome and characterized by tachycardia, profuse sweating, increased bronchial secretions, increased body temperature, and restlessness. These symptoms resolved upon reduction of the dose.

11 DESCRIPTION

CARBAGLU tablets for oral suspension contain 200 mg of carglumic acid. Carglumic acid, the active substance, is a carbamoyl phosphate synthetase 1 (CPS 1) activator and is soluble in boiling water, slightly soluble in cold water, and practically insoluble in organic solvents.

Chemically, carglumic acid is N-carbamoyl-L-glutamic acid or (2S)-2-(carbamoylamino) pentanedioic acid, with a molecular weight of 190.16.

The structural formula is:



Molecular Formula: C₆H₁₀N₂O₅

The inactive ingredients of CARBAGLU are croscarmellose sodium, hypromellose, microcrystalline cellulose, silica colloidal anhydrous, sodium lauryl sulfate, and sodium stearyl fumarate.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Carglumic acid is a synthetic structural analogue of N-acetylglutamate (NAG) which is produced from glutamate and acetyl-CoA in a reaction catalyzed by N-acetylglutamate synthase (NAGS), a mitochondrial liver enzyme. NAG acts as the essential allosteric activator of carbamoyl phosphate synthetase 1 (CPS 1), a mitochondrial liver enzyme which catalyzes the first reaction of the urea cycle. The urea cycle, whose role is the disposition of ammonia, includes a series of biochemical reactions in the liver resulting in the conversion of ammonia into urea, which is then excreted through the urine. Carglumic acid acts as a CPS1 activator, improves or restores the function of the urea cycle, and facilitates ammonia detoxification and urea production.

12.2 Pharmacodynamics

In a retrospective review of the clinical course in 23 patients with NAGS deficiency, carglumic acid reduced plasma ammonia levels within 24 hours when administered with and without concomitant ammonia lowering therapies. No dose-response relationship has been identified.

Cardiac Electrophysiology

The effect of carglumic acid was evaluated in a Phase 1, randomized study in 76 healthy volunteers. The study suggests a lack of clinically relevant QT prolongation effect at the highest therapeutic dose level (250 mg/kg/day).

12.3 Pharmacokinetics

The pharmacokinetics of carglumic acid in healthy subjects following an intravenous (IV) infusion over 2 hours at 8 mg/kg or an oral administration at 100 mg/kg are summarized in Table 3.

Table 3: Mean (SD) Pharmacokinetic Parameter Values of Carglumic Acid in Healthy Subjects

PK parameter	IV infusion 8 mg/kg (N=10)	Oral 100 mg/kg (N=12)
C _{max} (ng/mL)	8613 (558)	3284 (321)
T _{max} (hr)#	2 (1-2)	3 (2-4)
AUC (ng*hr/mL)	24501 (1613)	31426 (2150)
T _{1/2} (hr)	31 (3)	25 (2)
CL (L/hr/kg)	0.34 (0.02)	N/A
Vd (L/kg)	15 (1)	N/A

#Median (range); N/A, not applicable

Absorption

Following an oral administration of CARBAGLU 100 mg/kg in healthy subjects, the absolute bioavailability was approximately 10%.

Distribution

Carglumic acid is not bound to plasma proteins.

Elimination

Carglumic acid is predominantly excreted by the kidneys as unchanged product.

Metabolism

A proportion of carglumic acid may be metabolized by the intestinal bacterial flora.

The likely end product of carglumic acid metabolism is carbon dioxide, eliminated through the lungs.

Excretion

Following an oral administration of radiolabeled CARBAGLU at 100 mg/kg, 9% of the dose is excreted unchanged in the urine and up to 60% of the dose is recovered unchanged in the feces.

Specific Populations

Patients with Renal Impairment

The pharmacokinetics of carglumic acid in subjects with renal impairment were compared with healthy subjects with normal renal function following oral administration of a single dose of CARBAGLU 40 mg/kg or 80 mg/kg. The C_{max} and AUC_{0-t} of carglumic acid are

summarized in Table 4. The geometric mean ratio (90% CI) of AUC_{0-t} in subjects with mild, moderate, and severe renal impairment relative to those in their matched control subjects with normal renal function were approximately 1.3 (1.01, 1.68), 2.0 (1.65, 2.54), and 4.6 (3.36, 6.34) respectively [see *Dosage and Administration* (2.3)].

Drug Interaction Studies

Based on *in vitro* studies, carglumic acid is not an inducer of CYP1A1/2, CYP2B6, CYP2C, and CYP3A4/5 enzymes, and not an inhibitor of CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1, and CYP3A4/5 enzymes.

Based on *in vitro* studies, carglumic acid is a substrate of the human OAT1 transporter. Carglumic acid is not a substrate of MDR1, BCRP, MATE1, MATE2-K, OAT1, OAT3, OATP1B1, OATP1B3, OCT1, and OCT2. Carglumic acid is not an inhibitor of human BSEP, BCRP, MDR1, MATE1, MATE2-K, OAT1, OAT3, OATP1B1, OATP1B3, OCT1, and OCT2 transporters.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

The carcinogenic potential of carglumic acid was assessed in a 2-year carcinogenicity study in rats. Carglumic acid was not tumorigenic at oral doses up to 1000 mg/kg/day (approximately 34 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC).

Carglumic acid was negative in the Ames test, chromosomal aberration assay in human lymphocytes, and the *in vivo* micronucleus assay in rats.

There were no effects on fertility or reproductive performance in female rats at oral doses up to 2000 mg/kg/day (approximately 38 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC). In a separate study, mating and fertility were unaffected in male rats at oral doses up to 1000 mg/kg/day (approximately 34 times the maximum reported human maintenance dose [100 mg/kg/day] based on AUC).

14 CLINICAL STUDIES

14.1 Acute and Chronic Hyperammonemia due to NAGS Deficiency

The efficacy of CARBAGLU in the treatment of acute and chronic hyperammonemia due to NAGS deficiency was evaluated in a retrospective case series of 23 NAGS deficiency patients treated with CARBAGLU over a median duration of 7.9 years (range 0.6 to 20.8 years). For acute treatment, patients received CARBAGLU at 100 mg/kg/day to 250 mg/kg/day primarily administered in 2 to 4 divided doses. For maintenance treatment, the dosage was reduced over time based on plasma ammonia level and clinical response.

The baseline characteristics of the patient population are shown in Table 5.

Table 5: Baseline Characteristics of 23 NAGS Deficiency Patients Treated with CARBAGLU

Sex	Patients N=23	
	Male	14 (61%)
	Female	9 (39%)
Age at initiation of CARBAGLU therapy (years)	Mean (SD)	2 (4)
	Min-Max	0-13
Age groups at initiation of CARBAGLU therapy	<30 days	9 (39%)
	>30 days - 11 months	9 (39%)
	≥1 - 13 years	5 (22%)
NAGS gene mutations by DNA testing	Homozygous	14 (61%)
	Heterozygous	4 (17%)
	Not available	5 (22%)
Patients current treatment status	On-going	18 (78%)
	Discontinued	5 (22%)

Table 4: Mean (SD) C_{max} and AUC_{0-t} of Carglumic Acid Following Single Oral Dose Administration of CARBAGLU 80 mg/kg or 40 mg/kg in Subjects with Renal Impairment and Matched Healthy Control Subjects with Normal Renal Function

PK Parameters	Normal Renal Function 1a: eGFR ≥90 mL/min/1.73m ² (N=8)	Mild Renal Impairment: eGFR 60-89 mL/min/1.73m ² (N=8)	Moderate Renal Impairment: eGFR 30-59 mL/min/1.73m ² (N=8)	Normal Renal Function 1b: eGFR ≥90 mL/min/1.73m ² (N=8)	Severe Renal Impairment: eGFR ≤29 mL/min/1.73m ² (N=8)
	80 mg/kg			40 mg/kg	
C _{max} (ng/mL)	2983 (552)	4310 (1937)	6129 (1854)	1890 (901)	8377 (3815)
AUC _{0-t} (ng*hr/mL)	28313 (6204)	39545 (12109)	79766 (19708)	20212 (6186)	143075 (55910)

Treatment groups 1a and 1b represent two separate matched control groups of healthy subjects with normal renal function.

The clinical and biochemical data in the 23-patient case series were retrospectively collected, unblinded, and uncontrolled and preclude formal statistical testing. Short-term efficacy was evaluated using mean and median change in plasma ammonia levels from baseline to days 1 to 3. Persistence of the effect was evaluated using long-term mean and median change in plasma ammonia level. Of the 23 NAGS deficiency patients in the case series, 13 patients had documented plasma ammonia levels prior to CARBAGLU treatment and after long-term treatment with CARBAGLU and were evaluable. Table 5 summarizes the plasma ammonia levels at baseline, days 1 to 3 post-CARBAGLU treatment, and long-term CARBAGLU treatment (mean 8 years) in the 13 evaluable patients.


All 13 patients had increased plasma ammonia levels at baseline (mean 271 micromol/L; normal range: 5 to 50 micromol/L). By day 3 and with long-term treatment, normal plasma ammonia levels were attained (Table 6).

Table 6: Plasma Ammonia Levels in NAGS Deficiency Patients at Baseline and After Treatment with CARBAGLU

Timepoint	Patients (N = 13)	Ammonia level (micromol/L)
Baseline (prior to first dose of CARBAGLU)	N	13
	Mean (SD)	271 (359)
	Median	157
	Range	72-1428
	Missing Data	0
Day 1	N	10
	Mean (SD)	181 (358)
	Median	65
	Range	25-1190
	Missing Data	3
Day 2	N	8
	Mean (SD)	69 (78)
	Median	44
	Range	11-255
	Missing Data	5
Day 3	N	5
	Mean (SD)	27 (11)
	Median	25
	Range	12-42
	Missing Data	8
Long-term treatment (mean duration 8 years; median duration 6 years; range 1-16 years based on last available value while on CARBAGLU treatment)	N	13
	Mean (SD)	23 (7)
	Median	24
	Range	9-34
	Missing Data	0

The mean plasma ammonia level at baseline and the decline that is observed after treatment with CARBAGLU in 13 evaluable patients with NAGS deficiency is illustrated in Figure 1.

Figure 1: Mean Plasma Ammonia in 13 Evaluable NAGS Deficiency Patients at Baseline and After Treatment with CARBAGLU




14.2 Acute Hyperammonemia due to PA and MMA

A randomized, double-blind, placebo-controlled, multicenter clinical trial evaluated the efficacy of CARBAGLU in the treatment of hyperammonemia in patients with PA and MMA (NCT01599286).

Eligible hyperammonemic episodes, defined as an admission to the hospital with a plasma ammonia level ≥ 70 micromol/L, were randomized 1:1 to receive either CARBAGLU or placebo for 7 days or until hospital discharge, whichever occurred earlier. All patients received standard of care, including a combination of protein restriction, intravenous glucose, insulin, and/or L-carnitine; the use of alternative pathway medications (e.g., sodium benzoate and medications with phenylacetate as an active metabolite) was prohibited. CARBAGLU was dosed at 150 mg/kg/day for patients ≤ 15 kg or 3.3 g/m²/day for patients > 15 kg and was divided into 2 equal doses administered 12 hours apart by NG tube, G-tube, or oral syringe. Plasma ammonia testing was performed at pre-randomization and at post-dosing intervals of every 6-12 hours for the first 48 hours and every day thereafter if the ammonia level was ≥ 50 micromol/L.

The efficacy evaluation was based on 90 hyperammonemic episodes (42 treated with CARBAGLU and 48 with placebo) in 24 patients (12 male and 12 female) with PA (n = 15) or MMA (n = 9). The median patient age was 8 years (range 4 days to 29 years). The primary endpoint was the time from the first dose of drug to the earlier of plasma ammonia level ≤ 50 micromol/L (normal range) or hospital discharge. The median time to reach the primary endpoint was 1.5 days in the CARBAGLU group compared to 2.0 days in the placebo group, a difference of 0.5 days (95% confidence interval: -1.2, 0.1), driven exclusively by an effect on plasma ammonia normalization. Throughout the first three days of treatment, a higher proportion of CARBAGLU-treated episodes reached the primary endpoint compared to placebo-treated episodes (Figure 2).

Figure 2: Episodes Reaching the Earlier of Plasma Ammonia Level ≤ 50 micromol/L or Hospital Discharge in Patients with PA or MMA Treated with CARBAGLU or Placebo for up to 7 days



16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

CARBAGLU is a white and elongated 200 mg tablet for oral suspension, functionally scored with 3 lines for splitting into 4 equal portions, and coded "C" on one side.

CARBAGLU is available in 5 or 60 tablets in a high-density polyethylene bottle with child resistant polypropylene cap and desiccant unit.

NDC 52276-312-05 Bottles of 5 tablets

NDC 52276-312-60 Bottles of 60 tablets

Storage

Store in the original unopened container at 2°C to 8°C (36°F to 46°F).

After first opening of the container:

- Do not refrigerate, store at room temperature between 15°C and 30°C (59°F and 86°F).
- Keep the container tightly closed between openings in order to protect from moisture.
- Write the date of opening on the tablet container.
- Do not use after the expiration date stated on the tablet container.
- Discard one month after first opening.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Instructions for Use).

Instruct the patient on the following:

Preparation and Administration [see Dosage and Administration (2.3)]

- Disperse CARBAGLU tablets in water. Do not swallow whole or crushed.
- Take CARBAGLU immediately before meals or feedings.
- CARBAGLU tablets dispersed in water can be administered orally or via a nasogastric tube or gastrostomy tube as described in the *Instructions for Use*.

Storage [see How Supplied/Storage and Handling (16)]

- Store UNOPENED container in a refrigerator at 2°C to 8°C (36°F to 46°F).
- After first opening of the container: do not refrigerate, store at room temperature between 15°C and 30°C (59°F and 86°F). Keep the container tightly closed in order to protect from moisture. Write the date of opening on the tablet container.
- Discard one month after first opening. Do not use after the expiration date stated on the tablet container.

Pregnancy [see Use in Specific Populations (8.1)]

Advise women with NAGS deficiency who are exposed to CARBAGLU during pregnancy that there is a pregnancy surveillance program that monitors pregnancy outcomes.

Supplied by:
Recordati Rare Diseases
Puteaux, France

Licensed to and Distributed by:
Recordati Rare Diseases Inc.
Lebanon, NJ 08833



Carbaglu® is a licensed trademark of Recordati Rare Diseases Inc.

This product labeling may have been updated. For the most recent prescribing information, please visit www.recordatirarediseases.com/us or www.carbaglu.com.

INSTRUCTIONS FOR USE
CARBAGLU® (CAR-buh-gloo)
(carglumic acid)
Tablets for oral suspension

Important information:

- **CARBAGLU tablets for oral suspension (CARBAGLU tablet) must be mixed in water before taking.** CARBAGLU tablets should not be mixed in any other food or liquid.
- **Do not** swallow CARBAGLU tablets whole.
- **Do not** crush CARBAGLU tablets.
- Take CARBAGLU **right before** meals or feedings.
- The CARBAGLU tablet and water mixture has a slightly sour taste.

You may need to ask your healthcare provider or pharmacist for a medicine cup to measure the correct amount of water you will need to prepare the dose of CARBAGLU.

The CARBAGLU tablet has 3 lines used for splitting the tablet into 4 equal parts in order to get the prescribed dose. Ask your healthcare provider if you have any questions about how to split the tablet the right way or have any questions about the prescribed dose.

Taking CARBAGLU tablets by mouth using a cup:

Children and Adults

1. Add a minimum of 2.5 mL of water into a small cup for each CARBAGLU tablet, or each ½ or ¼ CARBAGLU tablet, needed for the prescribed dose. For example:
 - If the prescribed dose is 2 CARBAGLU tablets, add a minimum of 5 mL of water into the cup.
 - If the prescribed dose is 2 and a ¼ CARBAGLU tablets, add a minimum of 7.5 mL of water into the cup.
 - If the prescribed dose is 2 and a ½ CARBAGLU tablets, add a minimum of 7.5 mL of water into the cup.
 - Ask your healthcare provider if you are not sure of how much water you should use for the prescribed dose of CARBAGLU.
2. Place the prescribed number of CARBAGLU tablets into the water in the cup.
3. Carefully stir the CARBAGLU tablet and water mixture in the cup to avoid spilling the mixture. CARBAGLU tablets do not dissolve completely in water.
4. Swallow the CARBAGLU tablet and water mixture **right away**.
5. Pieces of the tablet may remain in the cup. Add more water to the cup to rinse the cup and swallow the mixture **right away**.
6. Repeat step 5 until there are no pieces of the tablet left in the cup.

Taking CARBAGLU tablets by mouth using an oral syringe:

Children

1. Add a minimum of 2.5 mL of water into a small cup for each CARBAGLU tablet, or each ½ or ¼ CARBAGLU tablet, needed for the prescribed dose. For example:
 - If the prescribed dose is 2 CARBAGLU tablets, add a minimum of 5 mL of water into the cup.
 - If the prescribed dose is 2 and a ¼ CARBAGLU tablets, add a minimum of 7.5 mL of water into the cup.

- If the prescribed dose is 2 and a ½ CARBAGLU tablets, add a minimum of 7.5 mL of water into the cup.
 - Ask your healthcare provider if you are not sure of how much water you should use for the prescribed dose of CARBAGLU.
2. Place the prescribed number of CARBAGLU tablets into the water in the cup.
 3. Carefully stir the CARBAGLU tablet and water mixture in the cup to avoid spilling the mixture. CARBAGLU tablets do not dissolve completely in water.
 4. Draw up all of the CARBAGLU tablet and water mixture in the cup into an oral syringe.
 5. Give your child the CARBAGLU tablet and water mixture **right away** by placing the tip of the oral syringe along the inner cheek of their mouth, on either the right or left side. Slowly push all the way down on the plunger to give the medicine.
 6. Pieces of the tablet may remain in the oral syringe. Refill the oral syringe with a minimum of 1 mL to 2 mL of water, and give your child the mixture **right away**.
 7. Repeat step 6 until there are no pieces of the tablet left in the oral syringe.

Giving CARBAGLU tablets through a nasogastric (NG) or gastrostomy tube (G-tube):

Children and Adults

1. Add a minimum of 2.5 mL of water into a small cup for each CARBAGLU tablet, or each ½ or ¼ CARBAGLU tablet, needed for the prescribed dose. For example:
 - If the prescribed dose is 2 CARBAGLU tablets, add a minimum of 5 mL of water into the cup.
 - If the prescribed dose is 2 and a ¼ CARBAGLU tablets, add a minimum of 7.5 mL of water into the cup.
 - If the prescribed dose is 2 and a ½ CARBAGLU tablets, add a minimum of 7.5 mL of water into the cup.
 - Ask your healthcare provider if you are not sure of how much water you should use for the prescribed dose of CARBAGLU.
2. Place the prescribed number of CARBAGLU tablets into the water in the cup.
3. Carefully stir the CARBAGLU tablet and water mixture in the cup to avoid spilling the mixture. CARBAGLU tablets do not dissolve completely in water.
4. Draw up all of the CARBAGLU tablet and water mixture in the cup into a catheter-tip syringe.
5. Connect the catheter-tip syringe to the NG tube or G-tube.
6. Give the CARBAGLU tablet and water mixture through the NG tube or G-tube **right away**.
7. Pieces of the tablet may remain in the catheter-tip syringe or NG tube or G-tube.
8. Refill the catheter-tip syringe with 1 mL to 2 mL of water and flush the NG tube or G-tube **right away**.

9. Repeat step 8 until there are no pieces of the tablet left in the catheter-tip syringe or NG tube or G-tube.

How should I store CARBAGLU?

- **Before opening**, store CARBAGLU in a refrigerator between 36°F to 46°F (2°C to 8°C) in the container it comes in.
- **After opening**, store CARBAGLU at room temperature between 59°F to 86°F (15°C to 30°C). **Do not** store CARBAGLU in a refrigerator after opening.
 - Keep CARBAGLU tablets in a tightly closed container to protect the tablets from moisture.
 - Write the date the CARBAGLU tablet container is opened on the container label. Throw away any unused tablets one month after opening the tablet container.
- Do not use CARBAGLU tablets after the expiration date on the tablet container.
- **Keep CARBAGLU and all medicines out of the reach of children.**

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Distributed by:

Recordati Rare Diseases Inc.
Lebanon, NJ 08833



Issued: 8/2021

Carbaglu® is a licensed trademark of Recordati Rare Diseases

PP-CBGL-US-0309